

A nonperturbative determination of C_A and the scaling of f_π and $m^{\overline{MS}}$

UKQCD Collaboration, presented by S. Collins^{a*}

^aDept. of Physics and Astronomy, Glasgow University, Glasgow, G12 8QQ, Scotland

We calculate the $O(a)$ improvement coefficient for the axial-vector current using the nonperturbative method proposed by the LANL group. Results for the coefficient in the range $\beta = 5.93$ to 6.2 are presented. We find c_A is close to the 1-loop tadpole-improved perturbative value. In addition, scaling of the pseudoscalar decay constant and renormalised quark mass is improved compared to that obtained using the (larger) c_A values obtained by the ALPHA collaboration.

1. Introduction

Symanzik improvement offers reduced discretisation errors for masses and matrix elements at finite lattice spacing but requires the calculation of the coefficients which accompany the improvement terms. In the light quark regime these coefficients can be calculated nonperturbatively by imposing the chiral Ward identities on the lattice.

c_A is the $O(a)$ improvement coefficient for the axial-vector current.

$$A_4 \rightarrow A_4^I = A_4 + ac_A \partial_4 + O(a^2) \quad (1)$$

It has been calculated nonperturbatively by the ALPHA collaboration [1] and LANL group [2]. The results are summarized in figure 1. There is a significant discrepancy in the values of c_A at smaller β . This might be accounted for by the $O(a)$ ambiguity that exists in c_A when it is determined nonperturbatively. Nevertheless, larger values of c_A lead to stronger lattice spacing dependence of f_π and the quark mass, and increase the difficulty of performing a continuum extrapolation. In addition, the ALPHA results have a very different β dependence compared to the 1-loop perturbative values, which suggests they contain a large nonperturbative contribution. Our aim was to investigate how well c_A is determined using the LANL method, which only requires conventional analysis, rather than Schrödinger functional techniques of the ALPHA

collaboration.

c_A is determined nonperturbatively using the PCAC relation. In euclidean space, for zero momentum

$$\langle \partial_4 A_4(x) J \rangle = 2m_{PCAC} \langle P(x) J \rangle \quad (2)$$

where m_{PCAC} is the PCAC quark mass, $P = \bar{\psi} \gamma_5 \psi$ and J is any operator with the pseudoscalar quantum numbers. Defining $r_J(t) = \frac{\langle \partial_4 A_4(t) J \rangle}{\langle P(t) J \rangle}$, equation 2 becomes on the lattice

$$r_J(t) = m(t) = 2m_{PCAC} + O(a). \quad (3)$$

The size of the $O(a)$ terms depends on the number of states contributing to $r_J(t)$. At early times (t_{ex}), the significant contribution from excited states means the $O(a)$ term is larger than at later times when the ground state dominates (t_{gs}). The improvement of equation 1 removes the $O(a)$ terms for all states and hence the two masses

$$r_J(t_{gs}) + ac_A s_J(t_{gs}) = m_{imp}(t_{gs}) = 2m_{PCAC} + O(a^2). \quad (4)$$

$$r_J(t_{ex}) + ac_A s_J(t_{ex}) = m_{imp}(t_{ex}) = 2m_{PCAC} + O'(a^2). \quad (5)$$

where $s_J(t) = \frac{\langle \partial_4^2 P(t) J \rangle}{\langle P(t) J \rangle}$, are equal up to higher order terms. We can use these equations to solve for c_A :

$$-\frac{1}{a} \frac{r(t_{ex}) - r(t_{gs})}{s(t_{ex}) - s(t_{gs})} \equiv c_A \quad (6)$$

The LANL method involves performing a fit to $r_J(t) + ac_A s_J(t) = \text{constant} = 2m$ over the fitting

*In collaboration with C. Davies (Glasgow University), G. P. Lepage (Cornell University), and J. Shigemitsu (Ohio-State University)




Figure 1. Summary of c_A determinations. LANL-2 and LANL-3 refer to c_A determined using 2-point and 3-point lattice derivatives respectively, as described in reference [2].

range t_{ex} to t_{gs} and is numerically equivalent to equation 6; c_A and $2m$ are extracted from the fit. The advantage of performing a fit is that one can test the ansatz with the χ^2 .

The $O(a)$ ambiguity in c_A is due to $O(a^2)$ terms in the axial-vector current, the pseudoscalar current and the light quark action. We define c_A in the chiral limit, but at finite quark mass there is an additional source of $O(a)$ error if standard, symmetric, $O(a^2)$ lattice temporal derivatives are used. This error can be removed by using improved $O(a^4)$ lattice derivatives. While, both choices of lattice derivative lead to consistent values for c_A in the chiral limit, we found it advantageous to use improved derivatives as this weakened the dependence of c_A on the quark mass and made the chiral extrapolation easier. We also found a wider window in t_{ex} from which c_A can be extracted.

2. Results for c_A

We calculated c_A on the UKQCD Collaboration quenched data set, at $\beta = 5.93, 6.0$ and 6.2 ; simulation details can be found in [3]. The best analysis was possible at $\beta = 5.93$.

Table 1 presents the results for c_A at $\beta = 5.93$ for “fuzzed” source [3] and LL correlators, where improved lattice derivatives have been used. We see that there is agreement between the results from different smearings and the results are stable

t_{ex}	$\kappa = 0.1327$	$\kappa = \kappa_c$
<i>FL</i>		
4	-0.037(2)	-0.050(3)
5	-0.035(4)	-0.052(8)
6	-0.029(9)	-0.022(19)
7	-0.025(15)	-
8	-0.066(264)	-
<i>LL</i>		
6	-0.024(6)	-0.032(14)
7	-0.022(11)	-
8	-0.046(30)	-

Table 1

c_A values extracted for $\beta = 5.93$ at finite quark mass and extrapolated to the chiral limit. For all results shown Q for the fit is > 0.01 . At κ_c and higher t_{ex} the statistical errors dominate and the results are not shown. $t_{gs} = 14$.

with t_{ex} . However, there is only a small window of timeslices for which Q for the fit is reasonable and before noise dominates. No significant change in c_A was found when t_{gs} was varied.

We take $c_A = -0.032(14)$ as our final value for this β . In choosing a value with a conservative error we aim to take into account the residual $O(a)$ ambiguity in c_A and some of the other systematic errors. In principle all results in the table are valid estimates of c_A . However, choosing a result with a much smaller error would require much more care in keeping the improvement conditions applied fixed when c_A is recalculated at other β s; thus ensuring, for example, any residual $O(a)$ error in c_A vanishes when the continuum limit is taken. In particular, with the LANL method one would need to keep the relative contribution of ground state to excited states at t_{ex} (in physical units) constant and this requires tuning of the smearing function, which was not possible in this study.

Repeating the analysis at $\beta = 6.0$ we found evidence of significant finite volume dependence. c_A decreased from $-0.039(6)$ at $\kappa = 0.13344$ on a 16^3 lattice to $-0.003(10)$ on a 32^3 lattice. The 16^3 lattice is 16% smaller in extent than that used at $\beta = 5.93$. Due to problems with chiral extrapolations, our final results as a function of t_{ex} either had too small a statistical error to re-




Figure 2. Scaling of the renormalised PS decay constant in units of r_0 .

flect the change in volume, or had too large an error to be useful. Hence, we disregard the results at this β in the later analysis. At $\beta = 6.2$, the lattice size is only 6% smaller, and we obtain $c_A = -0.031(5)$. We assume the latter error is sufficient to compensate for the small change in volume from $\beta = 6.2$ to 5.93.

3. Scaling of f_π and $m^{\overline{MS}}$

We calculate the renormalised pseudoscalar decay constant, given by

$$f^{ren} = Z_A(1 + ab_A m_q)(f^{(0)} + ac_A f^{(1)}) \quad (7)$$

$$f^{(0)} = \langle 0 | A_4 | P \rangle, f^{(1)} = \langle 0 | \partial_4 P | P \rangle \quad (8)$$

using our results for c_A and nonperturbative determinations of Z_A and the perturbative results for b_A ; f^{ren} is extracted at a reference mass $(r_0 M_{PS})^2 = 3.0$. This is compared in figure 2 with the results using the ALPHA collaboration values for c_A . We see that scaling is much better when our smaller values of c_A are used. The simultaneous fit to all 4 points only has a marginal $\chi^2 = 2.4$. More points and higher orders in a would be needed to include the ALPHA c_A values for f^{ren} .

The renormalised quark mass, at the same reference mass, was calculated using the method of reference [4]. The renormalisation-group invariant mass is computed from the PCAC mass using the nonperturbatively determined renormalisation factor Z_M : $\hat{M} = Z_M m_{PCAC}^{imp}$. The $\overline{MS}$ mass at 2 GeV is then obtained using 4-loop perturbation theory. Figure 3 shows that the scaling




Figure 3. Scaling of the renormalised quark mass in units of r_0 .

is better for smaller values of c_A ; the χ^2 for the combined fit is 3.3.

4. Conclusions

We implemented the LANL method for determining c_A in the range $\beta = 5.93$ to 6.2. Figure 1 shows our results are consistent with those of the LANL group. In particular, c_A does not change rapidly with β and is close to the 1-loop tadpole-improved value (using $\alpha_P(1/a)$). At $\beta = 6.0$, evidence of a finite volume dependence in c_A indicated care must be taken in keeping the physical volume in the simulation fixed when changing β with this method.

The scaling of both f_π and $m^{\overline{MS}}$ are improved using our, smaller, values of c_A compared to the values obtained by the ALPHA collaboration.

S. Collins acknowledges the support of a Royal Society of Edinburgh fellowship.

REFERENCES

1. ALPHA collaboration, M. Lüscher et. al., Nucl. Phys. B491 323 (1997).
2. T. Bhattacharya et. al., Phys. Lett. B461 79-88 (1999), T. Bhattacharya et. al., Phys. Rev. D63 074505 (2001). R. Gupta, these proceedings.
3. UKQCD collaboration, K. Bowler et. al, Phys. Rev. D62 054506 (2000), C. Allton et. al., hep-lat/0107021.
4. J. Garden et. al., Nucl. Phys. B571, 237-256(2000).